



4. D.U. Isamuxammedova “Shahar va Qishloq rekonstruksiylari” Toshkent-2018
5. S.M.Turobjonov, M.M.Niyazova, T.T.Tursunov, “Sanoat Chiqindilarini Rekuperatsiya qilish texnologiyasi” Toshkent 2011
6. O’rmonjonov Azizbek Muhammadjon o’g’li. (2025). INKLYUZIV TA’LIMNING MOHIYATI, MAQSADI VA RIVOJLANISH TENDENSIYALARI. (2025). Kelajak o’qituvchisi - Kelajak Bunyodkori, 1(2), 701-710.
7. Tursunov Ramziddin Abdixalil-o’g’li. (2025). JAMIYAT HAYOTIDA IJTIMOIIY ONG VA MA’NAVIYAT UYG’UNLIGI. TA’LIM VA TARAQQIYOT, 4(4), 324-330.

CLINICAL AND EPIDEMIOLOGICAL CHARACTERISTICS OF MEASLES IN THE SAMARKAND REGION

Rakhmonov Ravshan Nomozovich

Assistant of infectious diseases department of Samarkand state medical university

RESUME. Measles remains a significant public health challenge because of its extremely high contagiousness, ability to spread rapidly, and susceptibility of all age groups. International specialists emphasize that elimination of the measles virus is achievable through extensive vaccination programs. The primary contributing factors to the growing number of measles cases include gaps in vaccination coverage related to population migration, delayed identification of the disease, and the absence of timely anti-epidemic interventions.

Key words: measles, epidemiology, vaccination, morbidity, prevention.

SAMARQAND VILOYATIDA QIZAMIQ KASALLIGINING KLINIK VA EPIDEMIOLOGIK XUSUSIYATLARI

REZYUME. Qizamiq juda yuqori yuqumchangi, tez tarqalishi va barcha yosh guruhlari o’rtasida sezuvchanlik tufayli jiddiy jamoat salomatligi muammosi bo’lib qolmoqda. Xalqaro mutaxassislar qizamiq virusini butunlay yo’q qilish keng qamrovli emlash dasturlari orqali mumkinligini ta’kidlaydi. Qizamiq kasallanishining ortib borishiga asosiy omillar sifatida aholi migratsiyasi tufayli vaksinalash qamrovidagi bo’shliqlar, kasallikning kech aniqlanishi va o’z vaqtida epidemiya qarshi choralar ko’rilmaligi qayd etiladi.

Kalit so’zlar: qizamiq, epidemiologiya, emlash, kasallanish, profilaktika.

КЛИНИЧЕСКИЕ И ЭПИДЕМИОЛОГИЧЕСКИЕ ХАРАКТЕРИСТИКИ КОРИ В САМАРКАНДСКОЙ ОБЛАСТИ

РЕЗЮМЕ. Корь остаётся серьёзной проблемой общественного здравоохранения из-за её чрезвычайно высокой заразности, способности быстро распространяться и восприимчивости всех возрастных групп. Международные эксперты подчеркивают, что элиминация вируса кори достижима при условии широкого охвата вакцинацией. Основными факторами роста заболеваемости являются недостаточный уровень вакцинации вследствие миграционных процессов, позднее выявление заболевания и отсутствие своевременных противоэпидемических мероприятий.

Ключевые слова: корь, эпидемиология, вакцинация, заболеваемость, профилактика.

Relevance

Measles is an acute infectious viral illness characterized by respiratory involvement and systemic intoxication. The importance of the disease lies in its global prevalence, high transmissibility, and the fact that individuals of all ages can be affected. According to expert assessments, the eradication of measles is realistic due to vaccination. Over the past 15 years, the global mortality rate from measles has decreased by nearly 75% as a result of large-scale immunization activities. Despite this progress, the World Health Organization notes that outbreaks of measles infection continue to be registered in different parts of the world. Measles was officially declared eliminated in the United States in 2000, and a similar goal was set in Russia for 2010. Nevertheless, WHO reports that up to 3 million measles cases still occur globally every year. At present, outbreaks are documented in 14 European countries, and 9,500 cases were confirmed in



Europe in 2017. Measles can lead to serious complications involving the respiratory tract, gastrointestinal tract, and central nervous system. In some countries, measles remains an endemic disease. For example, 41,000 cases were reported in China in 2015, over 679,000 in India, and more than 10,000 cases in Pakistan, with children comprising the majority of those affected. It is noteworthy that 87% of infected individuals had never received measles vaccination. In Russia, measles incidence increased by 3.1% in 2017 compared with 2016. The rise in morbidity is attributed to vaccination refusal, delayed recognition of infection, and therefore delayed epidemiological interventions. Another contributing factor is insufficient vaccination coverage among certain population groups. Increased measles incidence is also associated with virus importation from other regions. The World Health Organization recognizes vaccine hesitancy as one of the major public health threats.

Purpose of the study. To determine the clinical and epidemiological characteristics of measles cases recorded in the Samarkand region during November and December of 2023.

Materials and methods. A total of 795 patients diagnosed with measles and hospitalized at the regional clinical infectious diseases hospital in Samarkand from November to December 2023 were observed. Of these patients, 716 were children between 1 and 3 years old. Diagnosis was based on epidemiological history, medical history, and assessment of clinical symptoms including the catarrhal period, rash stage, and pigmentation stage. Particular attention was given to the presence of Koplik’s spots, rash characteristics, rash sequence, and pigmentation changes. All patients underwent routine laboratory tests, including complete blood count, urinalysis, and stool examination. Serological testing for measles-specific IgM confirmed the diagnosis at admission. Chest radiographs were performed when necessary to identify complications. All children were evaluated by a pediatrician and, when needed, by a neurologist.

Results and discussion. Analysis revealed that among 795 measles-confirmed patients, infants under 1 year constituted 670 cases (84.3%), children aged 1–3 years accounted for 46 cases (5.8%), children aged 3–7 years accounted for 54 cases (6.8%), and adolescents and adults represented 25 cases (3.1%). Vaccination history demonstrated that the affected individuals were not vaccinated against measles. In 12.6% of cases, neurological disorders were the documented reason for medical exemption. All infants under one year were unvaccinated (84.3%), while individuals who received a single dose of vaccine constituted 3.1%. Contact tracing showed that 82% of children had contact exposure in medical institutions, 13% were exposed within families, and for 5% the source of exposure was unknown. Evaluation of disease severity revealed that 85% of patients had a moderate form, 10% had a severe form, and 5% had mild or modified measles. Most children (68%) were admitted during the rash stage, while 32% were hospitalized in the catarrhal stage. The clinical course was cyclic in most cases except for patients with modified measles. The incubation period lasted an average of 4–5 days. The catarrhal stage lasted 2–3 days in infants under one year and 3–5 days in older children. All patients had fever ranging from 37.5°C to 40°C; 68% had a temperature of 37.5–38.5°C and 32% had temperatures above 38.5°C. Children experienced decreased appetite, restlessness, and irritability, particularly those under 1 year (84.3%). Older children (6.8%) reported headache, weakness, coughing, and rhinorrhea. All patients demonstrated eyelid swelling, conjunctival hyperemia of varying intensity, and photosensitivity. Koplik’s spots, observed as small gray-white lesions with a hyperemic border on the buccal mucosa, were detected in 84% of patients. Oral mucosal swelling and hyperemia were present in most children, while 5% showed less pronounced catarrhal symptoms.

The rash lasted 3–4 days and appeared initially behind the ears and around the nose, spreading to the trunk and then covering the entire body by days 3–4. The rash involved both flexor and extensor surfaces and was abundant in 92% of patients; only 8% had limited eruptions. In 5% of patients, rash progression did not follow the typical stage sequence. Hemorrhagic spots appearing alongside papular lesions were detected in 4% of cases. Rash onset was accompanied by significant intoxication symptoms and fever of 38.5–39°C. In 97% of patients, the cough became barking in nature. Intestinal disorders manifested as frequent liquid stools were observed in 33% of infants. Complications detected during the rash stage indicated a complicated disease course: obstructive bronchitis occurred in 12% of patients, stenosing laryngotracheitis in 8%, and bronchopneumonia in 82%. In infants under one year, measles was accompanied by hypoplastic anemia. Pigmentation appeared by day 2–4, and was observed in 86% of patients. During the pigmentation stage, intoxication symptoms gradually subsided, and the general condition improved. Maculopapular rash in all patients eventually underwent light peeling. The total duration of the rash period ranged from 7 to 12 days.

Conclusions



1. Of the 795 measles cases, most affected individuals were young children, particularly infants under one year (84.3%).
2. Moderate severity of disease was the most common clinical manifestation (85% of cases), and 10% of cases were severe.
3. Complications were more frequently observed in younger children, including obstructive bronchitis (15%), stenosing laryngotracheitis (13%), and bronchopneumonia (82%).

Literature:

1. Мухаммадиев И. С., Рахмонов Р. Н., Ибрагимова М. Ф. Эффективность применения кларитромицина при пневмонии с атипичной этиологией у детей //Golden Brain. – 2024. – Т. 2. – №. 3. – С. 110-115.
2. Rakhmonov R. N., Zubaydullayev S. V. o ‘g ‘li, & Kardjavova, GA (2024) //ACUTE MYOCARDITIS IN CHILDREN ON THE BACKGROUND OF BRONCHO-PULMONARY DISEASES. GOLDEN BRAIN. – Т. 2. – №. 3. – С. 70-75.
3. Namozovich R. R. Mansurov Jasur Choriyor ugli, Sobirov Og‘abek Sobir ugli ugli, & Allanazarov Alisher Boymuratovich.(2024) //Acute Obstructive Bronchitis in Children: Main Etiological and Clinical Features. EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE. – Т. 4. – №. 2. – С. 98-100.
4. Rakhmonov R. N. et al. HEPATITIS C: THE CURRENT STATE OF THE PROBLEM //Web of Medicine: Journal of Medicine, Practice and Nursing. – 2025. – Т. 3. – №. 1. – С. 339-347.
5. Раббимова Н. и др. Математическое моделирование и прогнозирование заболеваемости кожным лейшманиозом в республике узбекистан //Журнал проблемы биологии и медицины. – 2017. – №. 1 (93). – С. 104-107.
6. Anvarovna Y. N. et al. Clinical and Epidemiological Characteristics of Shigellosis in Adults at the Contemporary Stage //Central Asian Journal of Medical and Natural Science. – 2021. – Т. 2. – №. 3. – С. 311-318.
7. Bakhronov J.J. ETIOLOGY, EPIDEMIOLOGY AND PATHOGENESIS OF VIRAL HEPATITIS B // Экономика и социум. 2025. №8 (135). URL: <https://cyberleninka.ru/article/n/etiology-epidemiology-and-pathogenesis-of-viral-hepatitis-b> (дата обращения: 10.11.2025).
8. Bakhronov J. J. ETIOLOGY, EPIDEMIOLOGY AND PATHOGENESIS OF VIRAL HEPATITIS DELTA // Экономика и социум. 2025. №6-2 (133). URL: <https://cyberleninka.ru/article/n/etiology-epidemiology-and-pathogenesis-of-viral-hepatitis-delta> (дата обращения: 10.11.2025).
9. Nematov H. A., Bakhronov J. J. CYTOMEGALOVIRUS INFECTION //Web of Medicine: Journal of Medicine, Practice and Nursing. – 2024. – Т. 2. – №. 12. – С. 335-344.
10. Zhasurovich B. Z. OCCULT HBV INFECTION //SHOKH LIBRARY. – 2025.
11. Zhasurovich B. Z. PREGNANCY AND VIRAL HEPATITIS B //SHOKH LIBRARY. – 2025.
12. Jakhongir B. et al. HEPATITIS C IN PREGNANT WOMEN AND NEWBORNS //Вестник магистратуры. – 2023. – №. 7 (142). – С. 10-12.
13. Rukhshonabonu M. et al. CLINICAL MANIFESTATION OF LENNOX-GASTAUT SYNDROME //Вестник магистратуры. – 2023. – №. 7 (142). – С. 19-21.
14. Bakhronov J. J. Kholmurodov Sh. F., Yakhyayeva Kh. D., Kuchkorov Sh. B. The role of artificial intelligence in modern medicine //Innovations in Technology and Science Education. – 2024. – Т. 2. – №. 7. – С. 464-467.

**PREVALENCE AND ANALYSIS OF TORCH INFECTIONS AMONG WOMEN OF
REPRODUCTIVE AGE IN THE SAMARKAND REGION**

Yarmukhamedova Makhbuba Kudratovna

Associate Professor of the Department of Infectious Diseases, Samarkand State Medical University
Iskhakova Kamila

Group 216 of Faculty of General Medicine , Samarkand State Medical University.

Annotation. This article evaluates the frequency of TORCH infections among women of reproductive age in the Samarkand region. Although no targeted epidemiological study was conducted to estimate regional prevalence, the work analyzes screening data obtained during routine laboratory testing. The results primarily reflect the serological immune response to the pathogens that form the TORCH group.